

Assignment #9

Due: Friday, April 28, 2004
TA: Christian Schunck
Office: 26-265
Email: chs@mit.edu
Office hour: April 27 @ 5:30-6:30 pm in 26-265

1. Magnitude of Raman scattering cross sections

- Find the spontaneous Raman cross section assuming $\langle i|\mathbf{D}|g\rangle = \langle i|\mathbf{D}|f\rangle = dea_0$ where $|g\rangle$ is the ground state from which the atom (molecule) is excited, $|i\rangle$ is the intermediate state, and $|f\rangle$ is the final state. Use $\epsilon_{ig} = \hbar\omega_{ig}/H$ (Hartree = e^2/a_0), and assume $\omega_{ig} \gg \omega$, where ω is the frequency of the exciting light.
- Estimate the cross section σ for light at 5000 \AA being Raman shifted by 300 cm^{-1} from molecules whose first electronic excitation energy is $\approx 1 \text{ Ry}$. Take $d = 1$.
- If you replace $|f\rangle$ by $|g\rangle$ (so the rate is from $|g\rangle$ to $|g\rangle$) show that you obtain the Rayleigh scattering cross section. (**Note:** Assume the oscillator strength to be 1.)

2. Saturation spectroscopy

This problem guides you through the concepts of saturation spectroscopy. This is one of the techniques to perform Doppler-free spectroscopy, i.e. to extract a narrow line (with the natural linewidth) in a gas with a broad velocity distribution. It nicely illustrates the combination of homogenous and inhomogeneous line broadening. Saturation spectroscopy is frequently used to lock lasers to atomic lines.

You should not get into nasty integrals for this problem. The drawn lineshapes should clearly show the basic features, but don't have to be exact.

a) Homogeneous broadening

Consider a dilute gas of density n composed of atoms with resonant frequency ω_0 and linewidth Γ . The gas is exposed to monochromatic light of frequency $\omega_L = \omega_0 + \delta$ and intensity $I = s I_{SAT}$ where I_{SAT} is the saturation intensity. Let us ignore the effects of the motion of the atoms, i.e. consider temperature $T = 0$. What are the densities of atoms in the ground state n_1 and the excited state n_2 , including the effect of saturation? What is the cross-section for absorption? The gas is in a box of length L along the direction of the incoming light. What fraction of the light is absorbed? (This is a small fraction, so don't worry about the effect of light attenuation on the saturation of the sample).

b) The Bennet hole

Now, let's endow these atoms with a mass m and a temperature T . Let the incoming light have a wavevector \mathbf{k}_L along the z-axis. What is the population density distribution in the ground state $n_1(v_z)$ as a function of v_z , the component of velocity in the z-direction? You should find that the light "burns a hole" (known as the Bennet hole) in the distribution of absorbers. What is the position of the hole? How do the width and depth (relative to the population for $s = 0$) vary with saturation parameter s ?

c) **Inhomogeneous broadening**

Consider that we sweep the frequency of the incident laser ω_L and measure the (small) absorption of the beam. Determine the fraction of the light absorbed as a function of s and δ and compare with its value at $s = 0$ (you don't need to solve the integral). For high temperatures ($k_L \bar{v}_z \gg \Gamma$), what is the width of the absorption line? Does saturation affect the width?

d) **Saturation spectroscopy**

To actually get some benefit from saturating the gas, we introduce a second laser beam.

- i) We can add a weak probe beam at frequency ω_p with wavevector \mathbf{k}_p . What is the absorption of this beam, including the effects of the saturating beam (\mathbf{k}_L, ω_L)? Again, just write the integral, and take the length of the box along \mathbf{k}_p to be L . Draw the absorption line shape, identifying the position and width of its features.
- ii) We can also just retroreflect the original beam. Draw the population distribution $n_1(v_z)$ for $\omega_L \neq \omega_0$ and $\omega_L = \omega_0$. Identifying the depth of the Bennett hole(s), draw the lineshape of absorption of the retroreflected beam (i.e. we scan ω_L). What is the width of the central feature (at $\delta = 0$)?