

Photon Recoil Momentum in Dispersive Media

Gretchen K. Campbell, Aaron E. Leanhardt,* Jongchul Mun, Micah Boyd, Erik W. Streed, Wolfgang Ketterle, and David E. Pritchard†

MIT-Harvard Center for Ultracold Atoms, Research Laboratory of Electronics and Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

(Received 31 January 2005; published 4 May 2005)

A systematic shift of the photon recoil momentum due to the index of refraction of a dilute gas of atoms has been observed. The recoil frequency was determined with a two-pulse light grating interferometer using near-resonant laser light. The results show that the recoil momentum of atoms caused by the absorption of a photon is $n\hbar k$, where n is the index of refraction of the gas and k is the vacuum wave vector of the photon. This systematic effect must be accounted for in high-precision atom interferometry with light gratings.

DOI: 10.1103/PhysRevLett.94.170403

PACS numbers: 03.75.Dg, 39.20.+q, 42.50.Ct

The momentum of a photon in a dispersive medium is of conceptual and practical importance [1–6]. When light enters a medium with an index of refraction n , the electromagnetic momentum is modified [3–5]. Momentum conservation requires then that the medium also has a mechanical momentum. When a pulse of light enters the medium, the particles in the medium are accelerated by the leading edge of the pulse and decelerated by the trailing edge [5]. As a result, no motion is left in the medium after the pulse has passed. When light is absorbed or reflected in the medium, the momentum transfer occurs in units of $\hbar k$ or $n\hbar k$, where k is the vacuum wave vector. An absorbing surface is equivalent to photons leaving the medium without reflection and would therefore receive a momentum of $\hbar k$ per incident photon. In contrast, as shown in Ref. [5], a reflecting surface within the medium will recoil with a momentum of $2n\hbar k$ per photon. In this case, the standing wave formed by the incident and reflected light pulse transfers momentum to the medium which remains even after the light pulse has left. This modification of the recoil momentum has so far been observed only for light being reflected from a mirror immersed in a liquid [7,8].

Recently, there have been discussions about what happens when an atom within an atomic cloud absorbs a photon. If one assumes that after absorbing the photon, no motion is left in the medium, then the recoil momentum should be $\hbar k$ [9]. The same conclusion is reached when one assumes a very dilute, dispersive medium with the absorbing atom localized in the vacuum space between the particles of the medium [10]. However, Ref. [6] argues that the atom will recoil with a momentum of $n\hbar k$, which requires particles in the medium to receive a backward momentum (for $n > 1$) due to the interaction of the oscillating dipole moments of the particles in the dispersive medium and the absorbing atom. So both for reflection by a mirror and absorption by an atom, a photon in a dispersive medium behaves as if it has a momentum of $n\hbar k$.

In this Letter, we examine this issue experimentally, showing that the atom recoils with momentum $n\hbar k$. This

has important consequences for atom interferometers using optical waves to manipulate atoms by the transfer of recoil momentum. High-precision measurements of the photon recoil are used to determine the fine-structure constant α [11–16]. Further improvements in the accuracy of photon recoil measurements, combined with the value of α derived from the $(g - 2)$ measurements for the electron and positron [17–19], would provide a fundamental test of QED. At low atomic densities, where atom interferometers usually operate, the index of the refraction effect is relatively small. However, the accuracy of the best photon recoil measurements is limited by the uncertainty in the correction to the photon recoil due to the index of refraction. Here we operate an atom interferometer with Bose-Einstein




FIG. 1. Kapitza-Dirac interferometer. The first pulse out-coupled a small fraction of atoms into the $|\pm 2n\hbar k\rangle$ momentum states. The outcoupled atoms moved within the initial condensate. After a variable delay τ , a second pulse was applied, and atoms outcoupled by the second pulse interfered with those outcoupled by the first pulse. The laser beam was applied perpendicular to the long axis of the condensate; the polarization, \vec{E} , was parallel to it and to the applied magnetic field bias, \vec{B} . The atoms were imaged after 38 ms of ballistic expansion. The field of view is $0.5 \text{ mm} \times 1.5 \text{ mm}$.

condensates, which have a much higher density than laser cooled atomic clouds, and observe how the index of refraction modifies the atomic recoil frequency $\hbar\omega_{\text{rec}} = \frac{\hbar^2 k^2}{2m}$, where m is the atomic mass.

The essential idea of our experiment is to measure the recoil frequency interferometrically using a two-pulse Ramsey interferometer. The two pulses are optical standing waves separated by a delay time τ (Fig. 1). The first pulse diffracts the atoms in a ^{87}Rb condensate into discrete momentum states. During the delay time τ the phase of each momentum state evolves at a different rate according to its recoil energy. The second pulse recombines the atoms with the initial condensate. The recombined components have differing phases leading to interference fringes that oscillate at the two-photon recoil frequency. By measuring the resulting frequency, ω as a function of the standing wave detuning from the atomic resonance, we found a distinctive dispersive shape for ω that fits the recoil momentum as $n\hbar k$.

The experiment was performed using an elongated ^{87}Rb Bose-Einstein condensate (BEC) created in a cloverleaf-type Ioffe-Pritchard magnetic trap previously described in Ref. [20]. The condensate, containing 1.5×10^6 atoms, was produced in the $|5^2S_{1/2}, F=1, m_F=-1\rangle$ state, and had a Thomas-Fermi radius of 8 (90) μm in the radial (axial) direction, and the magnetic trap had a radial (axial) trap frequency of 81 (7) Hz.

The BEC was illuminated with an optical standing wave created by a retroreflected, π -polarized laser beam. Losses in the retroreflected beam were negligible. The polarization of the beam was optimized by suppressing Rayleigh superradiance [20]. The laser was detuned from the $5^2S_{1/2}, F=1 \rightarrow 5^2P_{3/2}, F=1$ transition at $\lambda = 780$ nm, and had a linewidth γ much smaller than Γ , the natural linewidth of the transition. The intensity of the $5 \mu\text{s}$ long pulse was set to outcouple $\approx 5\%$ of the atoms into each of the $|\pm 2n\hbar k\rangle$ momentum states with no appreciable population in higher momentum states. This ensured that the density of the original condensate was nearly constant throughout the measurement. After a variable time τ , a second identical pulse was applied. The time between the first pulse and the shutoff of the magnetic trap was fixed at $600 \mu\text{s}$, which was less than a quarter of the radial trap period. The momentum distribution of the condensate was imaged after 38 ms of ballistic expansion, long enough for the momentum states to be resolved. The images were obtained using resonant absorption imaging after first optically pumping the atoms to the $5^2S_{1/2}, F=2$ state. To measure the effect of spontaneous light scattering from the standing wave, the density of the condensate (and associated mean-field shift) was determined after applying a single $5 \mu\text{s}$ pulse to the condensate, and then immediately releasing it from the magnetic trap. The number of atoms in the condensate was determined by integrating the optical density of the absorption image, which in turn was

calibrated by fitting the Thomas-Fermi radius of unperturbed condensates in time of flight [21].

The recoil frequency was found by fitting the oscillations in the fraction of atoms in the $|0n\hbar k\rangle$ momentum state as a function of the delay τ (Fig. 2) with a cosine function and a Gaussian envelope:

$$A \exp\left(-\frac{\tau^2}{\tau_c^2}\right) \cos(\omega\tau + \phi) + C. \quad (1)$$

The observation of up to ten oscillations provided a precise value of the recoil frequency. The origin of the damping time τ_c and of the offset C will be discussed later.

Figure 3 shows our measured values for $\omega/2\pi$ as a function of the detuning, $\Delta/2\pi$. The measured values for the frequency clearly follow the dispersive shape of the index of refraction. The variation in $\omega/2\pi$ as a function of the detuning was 2 kHz across the resonance, much larger than the statistical error on the frequency fits of less than 100 Hz. This conclusively shows that the momentum transferred to the atom when a photon is absorbed is $n\hbar k$.

We now discuss in more detail how the atoms interact with optical standing waves. For the short duration of the applied pulses ($5 \mu\text{s}$) we can assume that the atoms do not move during the pulse and ignore the kinetic energy of the atoms (Raman-Nath approximation). The interaction can then be described by the application of the ac Stark potential due to the standing wave $V(z) = \frac{\hbar\omega_k^2}{\Delta} \sin^2(nkz)$, where




FIG. 2. Interference fringes oscillating at the recoil frequency. (a) Absorption images for $\tau = 10\text{--}50 \mu\text{s}$. The detuning was $\Delta/2\pi = +520$ MHz. The field of view is $0.5 \text{ mm} \times 1.5 \text{ mm}$. (b) Fraction of atoms in the $|0n\hbar k\rangle$ momentum state as a function of τ . The fringes were fit using Eq. (1). The fitted frequency was $\omega = 2\pi \times 15\,627(39)$ Hz with decay constant $\tau_c = 461(25) \mu\text{s}$. The signal was normalized using the total atom number in all momentum states. The systematic scatter of the data from the fit indicates the reproducibility of the single shot measurements.




FIG. 3. Recoil frequency as a function of detuning, $\Delta/2\pi$, showing the dispersive effect of the index of refraction. The average density of the condensate for the solid points was $1.14(4) \times 10^{14} \text{ cm}^{-3}$, giving rise to a mean-field shift of 880 Hz. The shaded area gives the expected recoil frequency including the uncertainty in the density. The dashed line is at $\omega = 4\omega_{\text{rec}} + \rho U/\hbar$, the expected value without index of refraction effects. The dotted line is at $4\omega_{\text{rec}} = 15068 \text{ Hz}$, the two-photon vacuum recoil frequency. The data shown as open diamonds had increased spontaneous light scattering due to σ^\pm light contamination in the laser beam. The increased light scattering led to a lower initial density in the condensate, thus leading to a smaller mean-field shift. The σ^\pm contamination allowed $\Delta m_F = \pm 1$ transitions, thus for small detunings the proximity to the $|1, -1\rangle \rightarrow |0', 0\rangle$ transition located at $\Delta/2\pi = -72 \text{ MHz}$ resulted in higher spontaneous scattering rates. The open points have been scaled upward to correct for this lower density.

Δ is the detuning between the optical frequency and the atomic transition, and ω_R is the Rabi frequency. This equation is valid for large detuning, $\Delta^2 \gg \Gamma^2/4$. The short pulse limit, describing Kapitza-Dirac scattering, is valid for short interaction times $t_p \ll 1/\omega_{\text{rec}} \approx 40 \mu\text{s}$. The first pulse outcouples a fraction of atoms into the momentum states $|\pm 2\ell n\hbar k\rangle$, where the population in the ℓ th momentum state is given by $P_\ell = J_\ell^2(\theta)$ [22,23], where for a square pulse, $\theta = \frac{\omega_R t_p}{2\Delta}$, and J_ℓ is the ℓ th-order Bessel function of the first kind. For $\theta < 1$ a negligible fraction of atoms is diffracted into states with $\ell > 1$, and we can restrict our discussion to the $|\pm 2n\hbar k\rangle$ states. For our experimental parameters $\theta = 0.45$. During the delay time τ the phase of the $|\pm 2n\hbar k\rangle$ states evolves at a faster rate than the $|0n\hbar k\rangle$ state due to the recoil energy, $E_{\text{rec}} = 4n^2\hbar\omega_{\text{rec}}$; hence, the wave function evolves as

$$|\psi(\tau)\rangle = |\psi_0\rangle [J_1(\theta)|\pm 2n\hbar k\rangle e^{-i4n^2\omega_{\text{rec}}\tau} + J_0(\theta)|0n\hbar k\rangle]. \quad (2)$$

At $t = \tau$ a second pulse is applied that partially recombines the momentum states. After applying the two pulses, the probability of finding the atoms in the $|0n\hbar k\rangle$ state, $\rho_0 = |\langle\psi(\tau + t_p)|0n\hbar k\rangle|^2$, is given by

$$\rho_0 = J_0^4(\theta) + 4[J_0^2(\theta)J_1^2(\theta) + J_1^4(\theta)]\cos(4n^2\omega_{\text{rec}}\tau). \quad (3)$$

As a function of τ the density of the zero momentum peak oscillates at $4n^2\omega_{\text{rec}}$.

So far, we have ignored the motion of the atoms during the delay time τ . The amplitudes of the recombined components interfere only where they spatially overlap. After the first pulse, the atoms in the $|\pm 2n\hbar k\rangle$ states move with the recoil velocity ($v_{\text{rec}} \sim 12 \mu\text{m/ms}$). As the overlap between the recoiling atoms and those at rest decreases, the interference fringes decay. The overlap integral for this decay is approximated as a Gaussian with time constant, $\tau_c \approx 0.75R_{\text{TF}}/v_{\text{rec}}$, where R_{TF} is the Thomas-Fermi radius of the condensate [24].

The index of refraction for the condensate is derived from its macroscopic polarization P . For a two level system, $P = \chi\epsilon_0 E = i\rho \frac{\mu^2}{\hbar} \frac{E}{\Gamma - i\Delta}$, where χ is the atomic susceptibility, ϵ_0 is the permittivity of free space, μ is the dipole matrix element, ρ is the atomic density of the condensate. In this experiment the light was π polarized and detuned by Δ from the $5^2S_{1/2}, F=1 \rightarrow 5^2P_{3/2}, F'=1$ transition. For this polarization the selection rule is $\Delta m_F = 0$, and there are two allowed transitions from $|F=1, m_F=-1\rangle \rightarrow |1', -1\rangle$ and $|1, -1\rangle \rightarrow |2', -1\rangle$ that are separated by 157 MHz. Including both transitions in the derivation, the index of refraction, $n = \sqrt{1 + \chi}$, is given by

$$n = \sqrt{1 - 12\pi\rho\lambda^3 \left(\frac{5}{12} \frac{\Delta_1}{\Gamma} \frac{1}{1 + (\frac{\Delta_1}{\Gamma})^2} + \frac{1}{4} \frac{\Delta_2}{\Gamma} \frac{1}{1 + (\frac{\Delta_2}{\Gamma})^2} \right)}, \quad (4)$$

where Δ_1 and Δ_2 are the detunings relative to the $F=1 \rightarrow F'=1$ and $F=1 \rightarrow F'=2$ transitions, respectively. This equation is valid in the limit $\rho\lambda^3 \ll 1$ [25], where $\lambda = \lambda/2\pi$. For our experimental parameters $\rho\lambda^3 \approx 0.2$. In addition to the index of refraction shift, the observed recoil frequency has a mean-field shift [26]; the atoms in the $|\pm 2n\hbar k\rangle$ state have twice the mean-field energy of those at rest due to the exchange term in the interatomic potential. Including both the mean-field shift and the index of refraction, the frequency of the observed interference fringes should be

$$\omega = 4n^2\omega_{\text{rec}} + \frac{\rho U}{\hbar}, \quad (5)$$

where $\rho U = 4\pi\hbar^2 a \rho/m$, and a is the s -wave scattering length. The density $\rho = (4/7)\rho_0$, where ρ_0 is the peak condensate density and the factor of 4/7 is due to the inhomogeneous condensate density.

When the interference fringes were fit using Eq. (1), the average values for the amplitude A and offset C for all of the data points were 0.12(3) and 0.82(4), respectively. This is in reasonable agreement with the expected values of $A = 0.18$ and $C = 0.81$ for $\theta = 0.45$. For a Thomas-Fermi radius of $8 \mu\text{m}$ we would expect a decay time $\tau_c \approx 500 \mu\text{s}$. There was an unexplained shift in the fitted value

for τ_c between the red and blue side of the resonances: on the red side the average value was $347(20) \mu\text{s}$ and on the blue, $455(40) \mu\text{s}$.

The quadratic dependence in Eq. (5) on the index of refraction can be understood by considering the diffraction of atoms from the light grating created by the standing wave. When the first pulse is applied, the standing wave creates a grating with a periodicity $d = \lambda/2$. Within the condensate the index of refraction modifies the grating period by n , since $\lambda' = \lambda/n$. The momentum of atoms that diffract off the grating is changed by $\pm 2\hbar k'$, again within the condensate $k' = nk$, and the velocity of the atoms is modified. Assuming $n > 1$, when the second grating is applied the atoms have moved farther by a factor of n and the grating is shorter by n , changing the time scale for the interference fringes by a factor of n^2 .

The increase in the momentum transferred to the atoms can also be explained by considering the momentum transferred to atoms by a classical field. A derivation using the Lorentz force applied to the atoms during the absorption of a photon can be found in [6]. In a dielectric medium with $n > 1$, the magnetic field and therefore the Lorentz force are not modified. However, the electric field is weaker, and therefore it takes longer for the atom to perform half a Rabi cycle and be transferred to the excited state. During that longer time, the Lorentz force imparts a momentum to the atom which is larger than $\hbar k$.

For Kapitza-Dirac scattering, atoms are diffracted symmetrically into the $|\pm 2\ell n \hbar k\rangle$ momentum states, so momentum is clearly conserved. However, for processes such as Bragg scattering, where the atoms are scattered in only one direction, the index of refraction has an additional effect. Assuming a $\pi/2$ pulse with counterpropagating beams, where half the atoms are diffracted, for $n > 1$ the recoil momentum is a factor of n higher than in vacuum. For momentum to be conserved, the remaining atoms must recoil backwards with momentum $p = 2(n-1)\ell \hbar k$. For small fractional outcoupling the effect is negligible, since the extra momentum is distributed among the remaining condensate. However, if a large fraction of the condensate is outcoupled and ℓ is large, this effect could potentially be resolved in ballistic expansion.

We have discussed here the dispersive effect on the photon momentum near a one-photon resonance. An analogous effect occurs near two-photon resonances. In this case, the atomic polarizability is determined in third-order perturbation theory, and the resulting index of refraction has a sharp, narrow dispersive feature near the two-photon resonance [27]. In recent experiments at Stanford [13], such two-photon effects have been the leading source of uncertainty in high-precision determinations of atomic recoil frequencies and the fine-structure constant α .

In conclusion, we have measured a systematic shift in the photon recoil frequency due to the index of refraction of the condensate. This is the first direct observation of the

atomic recoil momentum in dispersive media. For high atomic densities, this shift can have a significant effect on atom interferometers, and is of particular importance for precision measurements of \hbar/m and α with cold atoms [13,16].

The authors thank A. Schirotzek, S. Chu, S. Harris, and H. Haus for insightful discussions, and M. Kellogg for experimental assistance. This work was supported by NSF and ARO.

*Present address: JILA, Boulder, CO 80309, USA.

†Electronic address: http://cua.mit.edu/ketterle_group/

- [1] H. Minkowski, *Nachr. Ges. Wiss. Göttingen* **53** (1908).
- [2] H. Minkowski, *Math. Ann.* **68**, 472 (1910).
- [3] M. Abraham, *Rend. Circ. Mat. Palermo* **28**, 1 (1909).
- [4] M. Abraham, *Rend. Circ. Mat. Palermo* **30**, 33 (1910).
- [5] J. P. Gordon, *Phys. Rev. A* **8**, 14 (1973).
- [6] M. P. Haugan and F. V. Kowalski, *Phys. Rev. A* **25**, 2102 (1982).
- [7] R. V. Jones and J. C. S. Richards, *Proc. R. Soc. London A* **221**, 480 (1954).
- [8] R. V. Jones and B. Leslie, *Proc. R. Soc. London A* **360**, 347 (1978).
- [9] J. M. Hensley, A. Wicht, B. C. Young, and S. Chu, in *Atomic Physics 17, Proceedings of the 17th International Conference on Atomic Physics*, edited by E. Arimondo, P. D. Natale, and M. Inguscio (AIP, Melville, NY, 2001).
- [10] This argument has been made previously by one of the authors (W. K.).
- [11] B. Taylor, *Metrologia* **31**, 181 (1994).
- [12] D. S. Weiss, B. C. Young, and S. Chu, *Phys. Rev. Lett.* **70**, 2706 (1993).
- [13] A. Wicht, J. M. Hensley, E. Sarajlic, and S. Chu, *Phys. Scr.* **T102**, 82 (2002).
- [14] S. Gupta, K. Dieckmann, Z. Hadzibabic, and D. E. Pritchard, *Phys. Rev. Lett.* **89**, 140401 (2002).
- [15] R. Battesti *et al.*, *Phys. Rev. Lett.* **92**, 253001 (2004).
- [16] Y. L. Coq *et al.*, *cond-mat/0501520*.
- [17] R. S. Van Dyck, P. B. Schwinberg, and H. G. Dehmelt, *Phys. Rev. Lett.* **59**, 26 (1987).
- [18] T. Kinoshita, *Phys. Rev. Lett.* **75**, 4728 (1995).
- [19] V. W. Hughes and T. Kinoshita, *Rev. Mod. Phys.* **71**, S133 (1999).
- [20] D. Schneble *et al.*, *Science* **300**, 475 (2003).
- [21] Y. Castin and R. Dum, *Phys. Rev. Lett.* **77**, 5315 (1996).
- [22] P. Meystre, *Atom Optics* (Springer-Verlag, New York, 2001).
- [23] S. Gupta, A. E. Leanhardt, A. D. Cronin, and D. E. Pritchard, *C.R. Acad. Sci. IV Phys.* **2**, 479 (2001).
- [24] M. Trippenbach *et al.*, *J. Phys. B* **33**, 47 (2000).
- [25] O. Morice, Y. Castin, and J. Dalibard, *Phys. Rev. A* **51**, 3896 (1995).
- [26] J. Stenger *et al.*, *Phys. Rev. Lett.* **82**, 4569 (1999).
- [27] A. Schirotzek, Diploma thesis, Universität Hamburg, 2004.